

IS A PROMINENT STERNITE RELATED TO MASS IN *CENTROBOLUS* COOK, 1897?

M. Ian Cooper*

University of South Africa, South Africa.

*Correspondence email, cm.i@aol.com, +27714620070.

Abstract- Three species of *Centrobolus* were identified (*C. fulgidus*, *C. inscriptus*, *C. ruber*) based on morphology and confirmed using Scanning Electron Microscopy (SEM) of gonopod structure. One set of linear measurements was made from the SEM micrographs: (1) prominence of the sternite. Mass in three species was gaged. Sternite prominence and mass were positively related ($r=0.71890876$, Z score= 2.02449752 , $n=8$, $p=0.02145942$). *C. inscriptus* had the highest sternite prominence (0.5 or 50%) and the highest mass (2.48 g) while *C. ruber* males had the lowest sternite prominence (0.25 or 25%) and the lowest mass (1.28 g). This supports the function of the sternite as an hypallometric device in sperm competition. Mechanical fit and stimulatory one-size-fits-all arguments were confirmed.

I. INTRODUCTION

The red millipede genus *Centrobolus* is well known for studies on sexual size dimorphism (SSD) and displays prolonged copulation durations for pairs of individuals of the species [4-9]. *Centrobolus* is distributed in temperate southern Africa with northern limits on the east coast of southern Africa at -17° latitude South (S) and southern limits at -35° latitude S. It consists of taxonomically important species with 12 species considered threatened and includes nine vulnerable and three endangered species [27]. It occurs in all the forests of the coastal belt from the Cape Peninsula to Beira in Mocambique [26]. Spirobolida has two pairs of legs modified into gonopods on the eighth and ninth diplosegments [28]. In *Centrobolus* the coleopods are the anterior gonopods of leg-pair eight and can be classed as paragonopods or peltogonopods because they are fused into a single plate-like structure and play a subsidiary role as inseminating devices while leg-pair nine are sperm-transferring [1]. The sternites (or stigma-carrying plates [29]) prevent lateral shifting (stabilizer) and stretch the vulva sac in a medial plane [3].

The genital morphology and mechanics of copulation were figured in three *Centrobolus* species [1, 2]. These are worm-like millipedes that have female-biased SSD [4-9, 12-19, 24]. From the

results, correlations between coleopod sternite prominence and mass were checked for correlations.

• II. MATERIALS AND METHODS

Three species of *Centrobolus* were identified based on morphology and confirmed using Scanning Electron Microscopy (SEM) of gonopod structure (*C. fulgidus*, *C. inscriptus*, *C. ruber*). The gonopods were dissected from males of these three species and prepared for SEM. Specimens were fixed, first in 2.5% glutaraldehyde (pH 7.4 phosphate-buffered saline) at 4°C for 24 hours, then in osmium tetroxide (2%). Dehydration through a graded alcohol series (50%, 60%, 70%, 80%, 90% to 100% ethanol) and critical point drying followed. Specimens were mounted on stubs and sputter-coated with gold palladium. Gonopods were viewed under a Cambridge S200 SEM. SEM micrographs were examined and the individual components of the gonopods were identified according to the available species descriptions. One set of linear measurements was made from the SEM micrographs: (1) prominence of the sternite (%). This has been estimated before as a ratio of how far it extends from the basal region up to the top of the coleopod. The collection of SEM micrographs for each species is particularly informative when comparisons are made between congruent views. These results have been published [1]. Dorsal tergite width was measured horizontally using Vernier calipers. Mass was gaged using a Mettler Autobalance. Sternite prominence and mass were correlated here using a Pearson Correlation Coefficient

(<https://www.gigacalculator.com/calculators/correlation-coefficient-calculator.php>).

Sternite prominence was correlated with mass in three

species (*C. fulgidus*, *C. inscriptus*, *C. ruber*) using a Pearson's Correlation Coefficient.

• III. RESULTS

Sternite prominence and mass were positively related ($r=0.71890876$, $Z \text{ score}=2.02449752$, $n=8$, $p=0.02145942$). Least-Squares Regression Line $y = 0.16553268 \cdot x + 0.07100380$. Sternite prominence and male mass were marginally related ($r=0.89067479$, $Z \text{ score}=1.42518105$, $n=4$, $p=0.07705248$). Sternite prominence and female mass were not related ($r=0.81915977$, $Z \text{ score}=1.15425805$, $n=4$, $p=0.12419725$).





Figure 1. Relationship between the prominence of the sternite (y) (%) and mass (g) (x) across three species of *Centrobolus* (*C. fulgidus*, *C. inscriptus*, *C. ruber*).



• IV. DISCUSSION

The genital morphology and mechanics of copulation were figured in three *Centrobolus* species [1, 2]. A direct relationship between an ultrastructural feature (sternite prominence) and mass of the millipedes is compared which certainly supports the function of the sternite as a device adapted for sperm competition [10, 30]. A relationship between this structural feature is present across three species suggesting adaptation to insemination. *C. inscriptus* had the highest sternite prominence (0.5 or 50%) and the highest mass (2.48 g) while *C. ruber* males had the lowest sternite prominence (0.25 or 25%) and the lowest mass (1.28 g). Although there was both a relationship between male nor female mass and sternite prominence, when mass is combined from both sexes a relationship exists. It can be difficult to understand the functionality and where there is no functional significance this could have been misjudged [25]. However, the sternites in *Centrobolus* millipedes predict a functional significance in assuring paternity. This is explained with mechanical fit and stimulatory one-size-fits-all arguments [22, 23].

• V. CONCLUSION

New relationships between ultrastructural features of the morphology (sternite prominence) and mass of the *Centrobolus* millipedes support the function

of the sternite as a hypoallometric device adapted toward reducing sperm competition and assuring paternity. A prominent sternite is hypoalloetrically related to body size.

• APPENDIX.

Male and female mass (g) and sternite prominence (%) in three species of *Centrobolus* with the first species (*C. inscriptus*) having two measurement sets.

- C. inscriptus* 2.48, 50 (male)
- C. inscriptus* 2.27, 50 (female)
- C. inscriptus* 2.00, 50 (male)
- C. inscriptus* 2.61, 50 (female)
- C. fulgidus* 1.29, 35 (male)
- C. fulgidus* 1.97, 35 (female)
- C. ruber* 1.28, 25 (male)
- C. ruber* 2.00, 25 (female)

• REFERENCES

- [1] M. I. Cooper, "Confirmation of four species of *Centrobolus* Cook (Spirobolida: Trigonulidae) based on gonopod ultrastructure," *Journal of Entomology and Zoology Studies*, vol. 4, no. 4, pp. 389-391, 2016.
- [2] M. I. Cooper, "Elaborate gonopods in the myriapod genus *Chersastus* (Diplopoda: Trigonulidae)," *Journal of Entomology and Zoology Studies*, vol. 3, no. 4, pp. 235-238, 2015.
- [3] M. Cooper, "Julid millipede and spirobolid millipede gonopod functional equivalents," *Journal of Entomology and Zoology Studies*, vol. 7, no. 4, pp. 333-335, 2019.
- [4] M. I. Cooper, "Sexual size dimorphism and corroboration of Rensch's rule in *Chersastus* millipedes," *Journal of Entomology and Zoology Studies*, vol. 2, no. 6, pp. 264-266, 2014.
- [5] M. I. Cooper, "Copulation and sexual size dimorphism in worm-like millipedes," *Journal of Entomology and Zoology Studies*, vol. 5, no. 3, pp. 1264-1266, 2017.
- [6] M. Cooper, "*Centrobolus anulatus* (Attems, 1934) reversed sexual size dimorphism," *Journal of Entomology and Zoology Studies*, vol. 6, no. 4, pp. 1569-1572, 2018.
- [7] M. I. Cooper, "The relative sexual size dimorphism of *Centrobolus inscriptus* compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 4, no. 6, pp. 504-505, 2016.
- [8] M. I. Cooper, "Relative sexual size dimorphism in *Centrobolus fulgidus* (Lawrence) compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 5, no. 3, pp. 77-79, 2017.
- [9] M. I. Cooper, "Relative sexual size dimorphism *Centrobolus ruber* (Attems) compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 5 no. 3, pp. 180-182, 2017.
- [10] M. I. Cooper, "Competition affected by re-mating interval in a myriapod," *Journal of Entomology and Zoology Studies*, vol. 3, no. 4, pp. 77-78, 2015.
- [11] M. Cooper, "Re-assessment of Rensch's rule in *Centrobolus*," *Journal of Entomology and Zoology Studies*, vol. 5, no. 6, pp. 2408-2410, 2017.
- [12] M. I. Cooper, "Size matters in myriapod copulation," *Journal of Entomology and Zoology Studies*, vol. 5, no. 2, pp. 207-208, 2017.
- [13] M. I. Cooper, "Sexual size dimorphism and the rejection of Rensch's rule in Diplopoda," *Journal of Entomology and Zoology Studies*, vol. 6, no. 1, pp. 1582-1587, 2018.
- [14] M. I. Cooper, "Allometry for sexual dimorphism in millipedes," *Journal of Entomology and Zoology Studies*, vol. 6, no. 1, pp. 91-96, 2018.
- [15] M. I. Cooper, "Trigonulid size dimorphism breaks Rensch," *Journal of Entomology and Zoology Studies*, vol. 6, no. 3, pp. 1232-1234, 2018.
- [16] M. Cooper, "A review of studies on the fire millipede genus *centrobolus* (diplopoda: trigonulidae)," *Journal of Entomology and Zoology Studies*, vol. 6, no. 4, pp. 126-129, 2018.
- [17] M. Cooper, "*Centrobolus sagatinus* sexual size dimorphism based on differences in horizontal tergite widths," *Journal of Entomology and Zoology Studies*, vol. 6, no. 6, pp. 275-277, 2018.
- [18] M. Cooper, "*Centrobolus silvanus* dimorphism based on tergite width," *Global Journal of Zoology*, vol. 3, no. 1, pp. 003-005, 2018.
- [19] M. Cooper, "Xylophagous millipede surface area to volume ratios are size dependent in forest," *Arthropods*, vol. 8, no. 4, pp. 127-136, 2019.
- [20] J. M. Dangerfield, S. R. Telford, "Seasonal activity patterns of julid millipedes in Zimbabwe," *Journal of Tropical Ecology*, vol. 7, pp. 281-285, 1991.
- [21] J. M. Dangerfield, A. E. Milner, R. Matthews, "Seasonal activity patterns and behaviour of juliform millipedes in south-eastern Botswana," *Journal of Tropical Ecology*, vol. 8, no. 4, pp. 451-464, 1992.
- [22] W. G. Eberhard, "STATIC ALLOMETRY AND ANIMAL GENITALIA," *Evolution*, vol. 63, no. 1, pp. 46-66, 2009.
- [23] W. G. Eberhard, B. A. Huber, R. L. Rodriguez Señor, R. D. Briceño, I. Salas, V. Rodriguez, "ONE SIZE FITS ALL? RELATIONSHIPS BETWEEN THE SIZE AND DEGREE OF VARIATION IN GENITALIA AND OTHER BODY PARTS IN TWENTY SPECIES OF INSECTS AND SPIDERS," *Evolution*, vol. 52, no. 2, pp. 415-431, 1998.
- [24] M. D. Greyling, R. J. Van Aarde, S. M. Ferreira, "Seasonal changes in habitat preferences of two closely related millipede species," *African Journal of Ecology*, vol. 39, no. 1, pp. 51-58, 2001.
- [25] G. I. Holwell, O. Kazakova, F. Evans, J. C. O'Hanlon, K. L. Barry, "The Functional Significance of Chiral Genitalia: Patterns of Asymmetry, Functional Morphology and Mating Success in the Praying Mantis *Ciulfina baldersoni*," *PLoS ONE*, vol. 10, no. 6, pp. e0128755, 2015.
- [26] R. F. Lawrence, "The Spirobolidae (Diplopoda) of the eastern half of Southern Africa*," *Annals of the Natal Museum*, vol. 18, no. 3, pp. 607-646, 1967.
- [27] R. P. Mailula, "Taxonomic revision and Red List assessment of the 'red millipede' genus *Centrobolus* (Spirobolida: Pachybolidae) of South Africa," *The University of Kwazulu natal*, xxiii+289, 2021.
- [28] P. Sierwald, J. E. Bond, "Current Status of the Myriapod Class Diplopoda (Millipedes): Taxonomic Diversity and Phylogeny,"

Annual Review of Entomology, vol. 52, no. 1, pp. 401-420, 2007.

- [29] T. Wesener, P. Sierwald, J-F. Wägele, "Sternites and spiracles – The unclear homology of ventral sclerites in the basal millipede order Glomeridesmida (Myriapoda, Diplopoda)," *Arthropod Structure & Development*, vol. 43, no. 1, pp. 87-95, 2014.
- [30] X. J. Zahnle, P. Sierwald, S. Ware, J. E. Bond, "Genital morphology and the mechanics of copulation in the millipede genus *Pseudopolydesmus* (Diplopoda: Polydesmida: Polydesmidae)," *Arthropod Structure & Development*, vol. 54, pp. 100913, 2020.