

AIR PRESSURE IS NOT RELATED TO SPECIES RICHNESS IN SPHAEROTHERIIDAE BRANDT, 1833

M. I. Cooper

University of Stellenbosch.

Abstract- Air pressure was tested for a relationship with species richness in pill millipedes *Sphaerotherium* and the outgroup species *Kylindotherium leve*. There were no differences between air pressure at 1 and 3 species (Z-test: P-value=0.208816, Zscore=0.810535, 3,3), 7 and 9 species (Z-test: P-value=0.277249, Z-score=0.591034; n=9, 7), between 3 and 7 species (Z-test: P-value=0.216748, Z-score=-0.783225; n=7, 3), or 24 species and 1 species (Z-test: P-value=0.441485, Z score=0.147205, n=24, 3), 24 species and 3 species (Z-test: P-value=0.142695, Z score=1.068290, n=24, 3), 24 species and 7 species (Z-test: P-value=0.403301, Z score=0.244813, n=24, 7), 24 species and 9 species (Z-test: P-value=0.128393, Z score=1.134022, n=24, 9), or 24 species and the rest (Z-test: P-value=0.130942, Z score=1.121951, n=24, 22). This suggested air pressure was not related to species richness in Sphaerotheriidae irrelevant of whether the outgroup *Kylindotherium leve* was included or excluded.

Keywords: air; diversity; gradient; latitude; pressure; richness; species.

I. INTRODUCTION

Species richness is the number of different species represented in an ecological community, landscape or region [4-7]. Species richness and biodiversity increase from the poles to the tropics for a wide variety of terrestrial and marine organisms and is referred to as a latitudinal diversity gradient (LDG) [17, 29]. Inverse LDG includes aphids, Chinese litter-dwelling thrips, diving beetle subfamily (Colymbetinae), European bryophytes, freshwater zooplankton, Holarctic tree frogs, ichneumonids, marine benthic algae, marine bivalves (Anomalodesmata), New World snake tribe (Lampropeltini), North American breeding birds, penguins, (peracarid) crustaceans, pitcher plant mosquito, pond turtles, Shallow-water molluscs, shorebirds, southeastern United States trees, subarctic forests and tropical leaf-litter ant communities [21, 22, 23, 25, 26, 28, 32, 33; 37, 38,42]. The LDG was measured and tested in the Oniscomorph forest millipede family Sphaerotheriidae [9]. This forest clade belonging to the Order Sphaerotheriida is distributed along the eastern coast of southern Africa consisting of species with concentrations around coastal bush

and forests [1, 2, 8, 15, 16, 19,24, 34, 39, 40, 41, 43]. Here species richness in pill millipedes are tested for correlations with air pressure in Sphaerotheriidae Brandt, 1833.

II. MATERIALS AND METHODS

46 of 49 valid species were identified as belonging to the genus *Sphaerotherium* Brandt, 1833 and the sister species *Kylindotherium leve* [16]. These were tabulated and known localities also listed [9]. Localities were obtained from literature [16]. GPS coordinates were obtained from internet sources for known localities using the locality followed with the keyword "GPS". Latitude and longitude coordinates were obtained. When co-ordinates were not in decimal degrees, they were subsequently converted to decimals in dividing the seconds in 60 and adding these to the minutes which were together divided through 60 to get the decimal behind or following the degree. Species accepted were in accordance with MilliBase (<http://www.millibase.org>). Species richness correlations with latitude were given [9]. Air pressure was calculated for each type locality at <https://www.mide.com/air-pressure-at-altitude-calculator>. P-value calculations were produced between air pressure at neighbouring species richness and between the highest species richness and the rest (Appendix 1). A test for normality of air pressure data was performed at <https://www.statskingdom.com/kolmogorov-smirnov-test-calculator.html>. The outcome of this test determined what P-value test would be used in comparing the data of air pressure across species richness. If the data were normal a T-test would be used while if the data were not normal a Z-test is used. The P-value calculator can be found at <https://www.gigacalculator.com/calculators/p-value-significance-calculator.php>.

III. RESULTS

There were no differences between air pressure at 1 and 3 species (Z-test: P-value=0.208816, Zscore=0.810535, 3,3), 7 and 9 species (Z-test: P-value=0.277249, Z-score=0.591034; n=9, 7), between 3 and 7 species (Z-test: P-value=0.216748, Z-score=-0.783225; n=7,6), or 24 species and 1 species (Z-test: P-value=0.441485, Z score=0.147205, n=24, 3), 24 species and 3 species (Z-test: P-value=0.142695, Z score=1.068290, n=24, 3), 24 species and 7 species (Z-test: P-value=0.403301, Z score=0.244813, n=24, 7), 24 species and 9 species (Z-test: P-value=0.128393, Z score=1.134022, n=24, 9), or 24 species and the rest (Z-test: P-value=0.130942, Z score=1.121951, n=24, 22). Results of the lilliefors test indicated that there is a significant difference from the normal distribution ($D(46) = 0.2$, $p = 0.00008$).

IV. DISCUSSION

Sphaerotherium and *Kylindotherium leve* are Gondwanan relicts [18]. *Sphaerotherium* are more temperate and show a general decline in LDG [5, 6, 7, 30, 44, 46]. Other groups showing an inverse LDG includes aphids, Chinese litter-dwelling thrips, diving beetle subfamily (Colymbetinae), European bryophytes, freshwater zooplankton, Holarctic tree frogs, ichneumonids, marine benthic algae, marine bivalves (Anomalodesmata), New World snake tribe (Lampropeltini), North American breeding birds, penguins, peracarid crustaceans, pitcher plant mosquito, pond turtles, Shallow-water molluscs, shorebirds, southeastern United States trees, subarctic forests and tropical leaf-litter ant communities [21, 22, 23, 25, 26, 28, 32, 33; 37, 38,42].

There may be an evolutionary preference for temperate environments appearing to have led to climatic constraints on dispersal based primarily on precipitation or temperature seasonality gradients [20, 32]. LDG depends on proximate factors affecting processes of speciation, extinction, immigration, and emigration and in

Sphaerotherium these factors are dependent on size which have been investigated in *Sphaerotherium* based on temperature, precipitation and latitude [10-12]. LDG may relate to body size in *Sphaerotherium* probably which does not agree with the trends in other taxa such as birds and fish [45]. The trend of a small body size associated with the inverse LDG is expected to be similar to the weak tendency found in mammals [14].

Although air pressure has been associated with species richness in red millipedes no relationship was found in the pill millipedes *Sphaerotherium* irrelevant of whether the outgroup *Kylindotherium* was included or excluded [13].

REFERENCES

- [1] Attems CMTG von. 1926. Myriopoda. In:Kükenthal-Krumbach. Handbuch der Zoologie, 4: 1-402
- [2] Attems CMTG von. 1928. The Myriopoda of SouthAfrica. Annals of the South African Museum, 26: 1-431
- [3] Brandt JF. 1833. Tentaminum quorundam monographicorum Insecta Myriapoda Chilognatha Latreillii spectantium prodromus. Bulletin de la Société Impériale des Naturalistes de Moscou, 6:194-209
- [4] Colwell RK. 2009. Biodiversity: Concepts, Patterns and Measurement In: The Princeton Guide to Ecology (Levin SA, ed), 257-263, Princeton, Princeton University Press, USA
- [5] Colwell RK, Hurtt GC. 1994. Non biological gradients in species richness and a spurious Rapoport effect. American Naturalist, 144: 570-595
- [6] Colwell RK, Lees DC. 2000. The mid-domain effect: geometric constraints on the geography of species richness. Trends in Ecology and Evolution, 15: 70-76
- [7] Colwell RK, Rahbek C, Gotelli NJ. 2004. The mid-domain effect and species richness patterns: what have we learned so far? American Naturalist, 163:E1-E23
- [8] Cooper MI. 2018. Sexual dimorphism in pill millipedes (Diplopoda). Journal of Entomology and Zoology Studies, 6(1): 613-616
- [9] Cooper M. Latitudinal gradient in species richness of *Sphaerotherium*. Arthropods, 2020, 9(4): 164-170
- [10] Cooper, M. I. Size across weather gradients in pill millipedes (Diplopoda): II. Female volume and

- lowest number of daily hours of sunshine (average). *Int. j. eng. sci. invention res. dev.* 2022; 9(2): 80-83.http://www.ijesird.com/aug_seven.PDF.
- [11] Cooper, M. I. Size across weather gradients in pill millipedes (Diplopoda): I. Female volume and precipitation. *Int. j. eng. sci. invention res. dev.* 2022;9(2): 84-87.
http://www.ijesird.com/aug_eight.PDF.
- [12] Cooper, M. Size across weather gradients in pill millipedes (Diplopoda): III. Female volume and lowest number of daily hours of sunshine (total), warmest month in the year, coolest month in the year, average annual temperature, and month with the highest number of rainy days. *Int. j. eng. sci. invention res. dev.* 2022; 9(2): 88-92.
http://www.ijesird.com/aug_nine.PDF.
- [13] Cooper, M. AIR PRESSURE IS RELATED TO SPECIES RICHNESS IN FOREST RED MILLIPEDES CENTROBOLUS COOK, 1897. *Int.j. eng. sci. invention res. dev.* 2023; 10(6): 1510-1534.
http://www.ijesird.com/december_7.pdf.
- [14] Gittleman JL, Purvis A. 1998. Body size and species-richness in carnivores and primates. *Proceedings of the Royal Society B: Biological Sciences*, 265(1391):113-119
- [15] Golovatch SI, Wesener T, Mauriès J-P, Semenyuk II. 2012. On the identities of *Cryxus* Leach, 1814 and *Zephronia* Gray, 1832, the oldest generic names in the millipede order Sphaerotheriida (Diplopoda). *Arthropoda Selecta*, 21 (4): 273-294
- [16] Hamer ML. 1998. Checklist of Southern African millipedes. *Annals of the Natal Museum*, 39(1): 39-43
- [17] Hillebrand H. 2004. On the Generality of the Latitudinal Diversity Gradient. *The American Naturalist*, 163(2): 192-211
- [18] Hopkin SP, Read HJ. 1992. Taxonomy, evolution, and zoogeography. In: *The Biology of millipedes*. Oxford University Press, UK
- [19] Jeekel CAW. 1951. A new pill-milliped from the Malayan Peninsula (Diplopoda, Sphaerotheriidae). *Tijdschrift voor Entomologie*, 93: 101-107
- [20] Kadamannaya BS, Sridhar KR, Sahadevan S. 2009. Seasonal Periodicity of Pill Millipedes (Arthrosphaera) and Earthworms of the Western Ghats, India. *World Journal of Zoology*, 4(2): 63-69
- [21] Kindlmann P, Dixon AFG, Traxmandlová-Schödelbauerová I. 2007. Inverse latitudinal gradients in species diversity. In: *Scaling Biodiversity* (Storch D, Marquet PA, ed). Cambridge University Press, Cambridge, UK
- [22] Krug AZ, Jablonski D, Valentine JW. 2007. Contrarian clade confirms the ubiquity of spatial origination patterns in the production of latitudinal diversity gradients. *Proceedings of the National Academy of Sciences of USA*, 104(46): 18129-18134
- [23] Kwon Y, Lee T, Lang A, Burnette D. 2019. Assessment on latitudinal tree species richness using environmental factors in the southeastern United States. *PeerJ*, 7: e6781
- [24] Lawrence RF. 1966. The Myriapoda of the Kruger National Park. *Zoologica Africana*, 2 (2): 225-262
- [25] Marshall KE, Baltzer JL. 2015. Decreased competitive interactions drive a reverse species richness latitudinal gradient in subarctic forests. *Ecology*, 96(2): 461-470
- [26] Mateo R, Broennimann O, Normand S, Petitpierre B, AraOyújo MB, Svenning J -C, Baselga A, FernándezGozález F, Rubio VG, Muñoz J. 2016. The mossy north: An inverse latitudinal diversity gradient in European bryophytes. *Scientific Reports*, 6: 25546
- [27] Mittelbach GC, Schemske GW, Cornell HV, Allen AP, Brown JM, Bush MB, Harrison SP, et al. 2007. Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters*, 10: 315-331
- [28] Morinière J, Van Dam MH, Hawlitschek O, Bergsten J, Michat MC, Hendrich L, Ribera I, Toussaint EFA, Balke M. 2016. Phylogenetic niche conservatism explains an inverse latitudinal diversity gradient in freshwater arthropods. *Scientific Reports*, 6: 26340
- [29] Pianka ER. 1989. Latitudinal Gradients in Species Diversity. *Trends in Ecology and Evolution*, 4(6):223
- [30] Pielou EC. 1977. The latitudinal spans of seaweed species and their patterns of overlap. *Journal of Biogeography*, 4: 299-311
- [31] Pitz KM, Sierwald P. 2010. Phylogeny of the millipede Order Spirobolida (Arthropoda: Diplopoda: Helminthomorpha). *Cladistics*, 26: 497-525
- [32] Pyron RA, Burbrink FT. 2009. Can the Tropical Conservatism Hypothesis explain temperate species richness patterns? An inverse latitudinal biodiversity gradient in the New World snake tribe Lampropeltini. *Global Ecology and Biogeography*, 18: 406-415
- [33] Rivadeneira MM, Thiel M, Gonzalez ER, Hays PA. 2011. An inverse latitudinal gradient of diversity of peracarid crustaceans along the Pacific Coast of South America: Out of the deep south. *Global Ecology and Biogeography*, 20(3): 437-448
- [34] Schubart O. 1958. Diplopoda II: Oniscomorpha. *South African Animal Life*, 5: 41-108

- [35] Sierwald P, Spelda J. 2020. MilliBase.<http://www.millibase.org>. Accessed on April 20, 2020
- [36] Sierwald P, Spelda J. 2018. MilliBase. *Sphaerotherium* Brandt, 1833. <http://www.millibase.org/aphia.php?p=taxdetails.A> ccessed on May 25, 2020
- [37] Silva RR, Brandão CRF. 2014. Ecosystem-wide morphological structure of leaf-litter ant communities along a tropical latitudinal gradient. PLoS ONE, 9(3):e93049
- [38] Sime KR, Brower AVZ. 1998. Explaining the latitudinal gradient anomaly in ichneumonid species richness: evidence from butterflies. Journal of Animal Ecology, 67: 387-399
- [39] Silvestri F. 1910. Materiali per una revisione dei Diplopoda Oniscomorpha. I. Specie del genere *Sphaerotherium* dell’Africa meridionale a me note. Bollettino del Laboratorio di Zoologia generale e agraria della R. Scuola superiore d’Agricoltura in Portici, 4: 180-220
- [40] Silvestri F. 1917. Materiali per una revisione dei Diplopoda Oniscomorpha. II. Specie di Sphaerotheridae delle regioni australiana e neozelandese a me note. Bollettino del Laboratorio di zoologia generale e agraria della Facoltà agraria in Portici, 12: 61-85
- [41] Van den Spiegel D, Golovatch SI, Hamer ML. 2002. Revision of some of the oldest species in the millipede genus *Sphaerotherium* Brandt, 1833 (Diplopoda, Sphaerotheriida, Sphaerotheriidae), with new synonymies. African Invertebrates, 43: 143-181
- [42] Wang J, Tong X, Donghui W. 2014. The effect of latitudinal gradient on the species diversity of Chinese litter-dwelling thrips. Zookeys (417): 9–20
- [43] Wesener T. 2016. The Giant Pill-Millipedes, order Sphaerotheriida - An annotated species catalogue with morphological atlas and list of apomorphies (Arthropoda: Diplopoda). Bonn Zoological Bulletin Supplementum, 63: 1-104
- [44] Willig MR, Lyons SK. 1998. An analytical model of latitudinal gradients of species richness with an empirical test for marsupials and bats in the New World. Oikos, 81: 93-98
- [45] Yen JDL, Thomson JR, Keith J, Paganin DM, Fleishman E, Bennett AF, Dobkin DS, Mac Nally R. 2018. Linking species richness and size diversity in birds and fishes. Ecography, 41(12): 1979-1991
- [46] Zapata FA, Gaston KJ, Chown SL. 2003. Mid-domain models of species richness gradients: assumptions, methods and evidence. Journal of Animal Ecology, 72: 677-690
- Appendix 1. Air pressure (Pa) followed by mean species richness in family Sphaerotheriidae genus *Sphaerotherium* and outgroup *Kylindotherium leve* (24).
 85556.60, 9
 86404.73, 7
 88987.01, 24
 101217.52, 24
 85556.60, 9
 91594.51, 24
 88987.01, 24
 100776.65, 7
 94406.69, 9
 94323.32, 7
 101217.52, 24
 101217.52, 24
 88987.01, 24
 82260.82, 7
 94421.16, 24
 101146.85, 25
 94406.69, 9
 99085.80, 24
 94550.04, 1
 98649.97, 7
 97895.21, 9
 100592.51, 1
 99035.51, 3
 84889.15, 3
 101002.63, 7
 89663.01, 1
 99597.56, 24
 101217.52, 24
 100697.17, 9
 85593.69, 24
 84337.14, 9
 88020.22, 3
 83274.52, 24
 99306.29, 7
 100775.23, 24
 93215.22, 24
 100758.33, 24
 101217.52, 24
 101217.52, 24
 89663.01, 9
 94841.70, 24
 100489.42, 24
 83274.52, 24
 88987.01, 24
 100564.98, 9
 99204.85, (24)